Factoring - Trinomials $\mathbf{a} = \mathbf{1}$

Factoring with three terms, or trinomials, is the most important type of factoring to be able to master. As factoring is multiplication backwards we will start with a multiplication problem and look at how we can reverse the process

Example 1.

$$\begin{array}{ll} (x+6)(x-4) & \text{Distribute}\,(x+6)\,\text{through second parenthesis} \\ x(x+6)-4(x+6) & \text{Distribute each monomial through parenthesis} \\ x^2+6x-4x-24 & \text{Combine like terms} \\ x^2+2x-24 & \text{Our Solution} \end{array}$$

You may notice that if you reverse the last three steps the process looks like grouping. This is because it is grouping! The GCF of the left two terms is x and the GCF of the second two terms is x. The way we will factor trinomials is make them into a polynomial with four terms and then factor by grouping. This is shown in the following example, the same problem worked backwards

Example 2.

$$x^2 + 2x - 24$$
 Split middle term into $+6x - 4x$
 $x^2 + 6x - 4x - 24$ Grouping: GCF on left is x , on right is -4
 $x(x+6) - 4(x+6)$ ($x+6$) is the same, factor out this GCF
($x+6$)($x-4$) Our Solution

The trick to make these problems work is how we split the middle term. Why did we pick +6x - 4x and not +5x - 3x? The reason is because 6x - 4x is the only combination that works! So how do we know what is the one combination that works? To find the correct way to split the middle term we will use what is called the ac method. In the next lesson we will discuss why it is called the ac method. The way the ac method works is we find a pair of numers that multiply to a certain number and add to another number. Here we will try to multiply to get the last term and add to get the coefficient of the middle term. In the previous example that would mean we wanted to multiply to -24 and add to +2. The only number that do this are 6 and -4 (6 $\cdot -4 = -24$ and 6 + (-4) = 2). This process is shown in the next few examples

Example 3.

$$x^2 + 9x + 18$$
 Want to multiply to 18, add to 9
 $x^2 + 6x + 3x + 18$ 6 and 3, split the middle term
 $x(x+6) + 3(x+6)$ Factor by grouping
 $(x+6)(x+3)$ Our Solution

Example 4.

$$x^2-4x+3$$
 Want to multiply to 3, add to -4 $x^2-3x-x+3$ -3 and -1 , split the middle term $x(x-3)-1(x-3)$ Factor by grouping

$$(x-3)(x-1)$$
 Our Solution

Example 5.

$$x^2 - 8x - 20$$
 Want to multiply to -20 , add to -8
 $x^2 - 10x + 2x - 20$ -10 and 2, split the middle term
 $x(x-10) + 2(x-10)$ Factor by grouping
 $(x-10)(x+2)$ Our Solution

Often when factoring we have two variables. These problems solve just like problems with one variable, using the coefficients to decide how to split the middle term

Example 6.

$$a^2 - 9ab + 14b^2$$
 Want to multiply to 14, add to -9
 $a^2 + 7ab + 2ab + 14b^2$ 7 and 2, splif the middle term
 $a(a+7b) + 2b(a+7b)$ Factor by grouping
 $(a+7b)(a+2b)$ Our Solution

As the past few examples illustrate, it is very important to be aware of negatives as we find the pair of numbers we will use to split the middle term. Consier the following example, done incorrectly, ignoring negative signs

Warning 7.

$$x^2 + 5x - 6$$
 Want to multiply to 6, add 5
 $x^2 + 2x + 3x - 6$ 2 and 3, split the middle term
 $x(x+2) + 3(x-2)$ Factor by gouping
??? Binomials do not match!

Because we did not use the negative sign with the six to find our pair of numbers, the binomials did not match and grouping was not able to work at the end. Now the problem will be done correctly.

Example 8.

$$x^2 + 5x - 6$$
 Want to multiply to -6 , add to 5
 $x^2 + 6x - x - 6$ 6 and -1 , splif the middle term
 $x(x+6) - 1(x+6)$ Factor by grouping
 $(x+6)(x-1)$ Our Solution

You may have noticed a shortcut for factoring these problems. Once we identify the two numbers that are used to split the middle term, these are the two numbers in our factors! In the previous example, the numbers used to split the middle term were 6 and -1, our factors turned out to be (x + 6)(x - 1). This pattern

does not always work, so becareful getting in the habit of using it. We can use it however, when we have no number in front of x^2 . In all the problems we have factored in this lesson there is no number in front of x^2 . If this is the case then we can use this shortcut. This is shown in the next few examples.

Example 9.

$$x^2 - 7x - 18$$
 Want to multiply to -18 , add to -7 -9 and 2, write the factors $(x-9)(x+2)$ Our Solution

Example 10.

$$m^2 - mn - 30n^2$$
 Want to multiply to -30 , add to -1
5 and -6 , write the factors, don't forget second variable $(m+5n)(m-6n)$ Our Solution

It is possible to have a problem that does not factor. If there is not combination of numbers that multiplies and adds to the correct numbers, then we say we cannot factor the polynomial, or we say the polynomial is prime. This is shown in the following example.

Example 11.

$$x^2 + 2x + 6$$
 Want to multiply to 6, add to 2
 $1 \cdot 6$ and $2 \cdot 3$ Only possibilities to multiply to six, none add to 2
Prime, can't factor Our Solution

When factoring it is important not to forget about the GCF. If all the terms in a problem have a common factor we will want to first factor out the GCF before we factor using any other method.

Example 12.

$$3x^2 - 24x + 45$$
 GCF of all terms is 3, factor this out $3(x^2 - 8x + 15)$ Want to multiply to 15, add to -8 -5 and -3 , write the factors $3(x-5)(x-3)$ Our Solution

Again it is important to comment on the shortcut of jumping right to the factors, this only works if there is no coefficient on x^2 . In the next lesson we will look at how this process changes slightly when we have a number in front of x^2 . Be careful not to use this shortcut on all factoring problems!



Beginning and Intermediate Algebra by Tyler Wallace is licensed under a Creative Commons Attribution 3.0 Unported License. (http://creativecommons.org/licenses/by/3.0/)

Practice - Trinomials where a=1

Factor each completely.

1)
$$p^2 + 17p + 72$$

3)
$$n^2 - 9n + 8$$

5)
$$x^2 - 9x - 10$$

7)
$$b^2 + 12b + 32$$

9)
$$x^2 + 3x - 70$$

11)
$$n^2 - 8n + 15$$

13)
$$p^2 + 15p + 54$$

15)
$$n^2 - 15n + 56$$

17)
$$u^2 - 8uv + 15v^2$$

19)
$$m^2 + 2mn - 8n^2$$

21)
$$x^2 - 11xy + 18y^2$$

23)
$$x^2 + xy - 12y^2$$

$$25) \ x^2 + 4xy - 12y^2$$

27)
$$5a^2 + 60a + 100$$

29)
$$6a^2 + 24a - 192$$

$$31) \ 6x^2 + 18xy + 12y^2$$

$$33) \ 6x^2 + 96xy + 378y^2$$

2)
$$x^2 + x - 72$$

4)
$$x^2 + x - 30$$

6)
$$x^2 + 13x + 40$$

8)
$$b^2 - 17b + 70$$

10)
$$x^2 + 3x - 18$$

12)
$$a^2 - 6a - 27$$

14)
$$p^2 + 7p - 30$$

16)
$$m^2 - 15mn + 50n^2$$

18)
$$m^2 - 3mn - 40n^2$$

20)
$$x^2 + 10xy + 16y^2$$

22)
$$u^2 - 9uv + 14v^2$$

24)
$$x^2 + 14xy + 45y^2$$

26)
$$4x^2 + 52x + 168$$

28)
$$5n^2 - 45n + 40$$

30)
$$5v^2 + 20v - 25$$

32)
$$5m^2 + 30mn - 90n^2$$

34)
$$6m^2 - 36mn - 162n^2$$



Beginning and Intermediate Algebra by Tyler Wallace is licensed under a Creative Commons Attribution 3.0 Unported License. (http://creativecommons.org/licenses/by/3.0/)

Answers - Trinomials wehre a=1

1)
$$(p+9)(p+8)$$

2)
$$(x-8)(x+9)$$

3)
$$(n-8)(n-1)$$

4)
$$(x-5)(x+6)$$

5)
$$(x+1)(x-10)$$

6)
$$(x+5)(x+8)$$

7)
$$(b+8)(b+4)$$

8)
$$(b-10)(b-7)$$

9)
$$(x-7)(x+10)$$

10)
$$(x-3)(x+6)$$

11)
$$(n-5)(n-3)$$

12)
$$(a+3)(a-9)$$

13)
$$(p+6)(p+9)$$

14)
$$(p+10)(p-3)$$

16)
$$(m-5n)(m-10n)$$

17)
$$(u-5v)(u-3v)$$

18)
$$(m+5n)(m-8n)$$

19) Not factorable

20)
$$(x+8y)(x+2y)$$

21)
$$(x-9y)(x-2y)$$

22)
$$(u-7v)(u-2v)$$

23)
$$(x-3y)(x+4y)$$

24)
$$(x+5y)(x+9y)$$

26)
$$4(x+7)(x+6)$$

27)
$$5(a+10)(a+2)$$

28)
$$5(n-8)(n-1)$$

29)
$$6(a-4)(a+8)$$

30)
$$5(v-1)(v+5)$$

31)
$$6(x+2y)(x+y)$$

32)
$$5(m^2+6mn-18n^2)$$

33)
$$6(x+9y)(x+7y)$$

34)
$$6(m-9n)(m+3n)$$



Beginning and Intermediate Algebra by Tyler Wallace is licensed under a Creative Commons Attribution 3.0 Unported License. (http://creativecommons.org/licenses/by/3.0/)